

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Estrogen Receptor, alpha (Marker of Estrogen Dependence) (rESR1/1935), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [rESR1/1935], Mouse, Monoclonal BOT-BNCB2300-100

| | |
|--------------------------|---|
| Artikelname | Estrogen Receptor, alpha (Marker of Estrogen Dependence) (rESR1/1935), Biotin conjugate, 0.1mg/mL, IgG1, Clone: [rESR1/1935], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB2300-100 |
| Hersteller Artikelnummer | BNCB2300-100 |
| Alternativnummer | BOT-BNCB2300-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human Estrogen Receptor alpha protein |
| Konjugation | Biotin |
| Produktbeschreibung | This MAb is specific to ER alpha and shows minimal cross-reaction with other members of the family. ER is an important regulator of growth and differentiation in the mammary gland. Presence of ER in breast tumors indicates an increased likelihood of ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [rESR1/1935] |
| Molekulargewicht | 67 kDa |
| Isotyp | IgG1 |
| UniProt | P03372 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 1-2 ug/mL Optimal dilution for a specific application should be determined by user |