

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cyclooxygenase-2 (COX-2)(COX2/1941), Biotin conjugate, 0.1mg/mL, Clone: [COX2/1941], Mouse, Monoclonal BOT-BNCB2377-500

| | |
|--------------------------|---|
| Artikelname | Cyclooxygenase-2 (COX-2)(COX2/1941), Biotin conjugate, 0.1mg/mL, Clone: [COX2/1941], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB2377-500 |
| Hersteller Artikelnummer | BNCB2377-500 |
| Alternativnummer | BOT-BNCB2377-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human COX2 protein fragment (around aa 442-572) (exact sequence is proprietary) |
| Konjugation | Biotin |
| Produktbeschreibung | Prostaglandins are a diverse group of autocrine and paracrine hormones that mediate many cellular and physiologic processes. Prostaglandin H2 (PGH2) is an intermediate molecule in formation of the prostaglandins. Cyclooxygenase-1 (Cox-1) and cyclooxy... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [COX2/1941] |

| | |
|------------------------|--|
| Molekulargewicht | 70-72 kDa |
| UniProt | P35354 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |