

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD25 / IL2RA (Activated Lymphocyte Marker)(IL2RA/2394), Biotin conjugate, 0.1mg/mL, Clone: [IL2RA/2394], Mouse, Monoclonal BOT-BNCB2394-500

| | |
|--------------------------|---|
| Artikelname | CD25 / IL2RA (Activated Lymphocyte Marker)(IL2RA/2394), Biotin conjugate, 0.1mg/mL, Clone: [IL2RA/2394], Mouse, Monoclonal |
| Artikelnummer | BOT-BNCB2394-500 |
| Hersteller Artikelnummer | BNCB2394-500 |
| Alternativnummer | BOT-BNCB2394-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human IL2RA protein fragment (around aa 42-183) (exact sequence is proprietary) |
| Konjugation | Biotin |
| Produktbeschreibung | Recognizes a protein of 55 kDa, identified as CD25. It is expressed on activated T- and B-cells and activated monocytes/macrophages. With respect to lymphomas, CD25 is present on malignant cells of Hodgkins disease, HTLV-1 adult T-cell leukemia, cuta... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [IL2RA/2394] |

| | |
|------------------------|--|
| Molekulargewicht | 55 kDa |
| UniProt | P01589 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |