

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **OLIG2 (Marker of Glial Brain Tumors) (OLIG2/2400), Biotin conjugate, 0.1mg/mL, Clone: [OLIG2/2400], Mouse, Monoclonal BOT-BNCB2400-100**

Artikelname	OLIG2 (Marker of Glial Brain Tumors) (OLIG2/2400), Biotin conjugate, 0.1mg/mL, Clone: [OLIG2/2400], Mouse, Monoclonal
Artikelnummer	BOT-BNCB2400-100
Hersteller Artikelnummer	BNCB2400-100
Alternativnummer	BOT-BNCB2400-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant fragment of human OLIG2 protein (around aa 1-141) (exact sequence is proprietary)
Konjugation	Biotin
Produktbeschreibung	Olig2, a basic helix-loop-helix transcription factor, is involved in oligodendroglial specification. Olig2 expression has been reported in most glial tumors, such as oligodendrogliomas and astrocytomas. Although more than half of glioblastomas are po...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[OLIG2/2400]
Molekulargewicht	30-40 kDa
UniProt	<a href="#">Q13516</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user