

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cyclin E (G1/S-Phase Cyclin) (CCNE1/2460), Biotin conjugate, 0.1mg/mL, Clone: [CCNE1/2460], Mouse, Monoclonal BOT-BNCB2460-500**

Artikelname	Cyclin E (G1/S-Phase Cyclin) (CCNE1/2460), Biotin conjugate, 0.1mg/mL, Clone: [CCNE1/2460], Mouse, Monoclonal
Artikelnummer	BOT-BNCB2460-500
Hersteller Artikelnummer	BNCB2460-500
Alternativnummer	BOT-BNCB2460-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human Cyclin E (CCNE1) protein fragment (around aa 10-176) (exact sequence is proprietary)
Konjugation	Biotin
Produktbeschreibung	Cyclin E belongs to the highly conserved cyclin family, whose members exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. Cyclins function as regulators of CDK kinases. Cyclin E fo...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[CCNE1/2460]
Molekulargewicht	53 kDa
UniProt	<a href="#">P24864</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user