

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Perforin-1 (Pore Forming Protein) (Apoptosis Marker)(PRF1/2467), Biotin conjugate, 0.1mg/mL, Clone: [PRF1/2467], Mouse, Monoclonal BOT-BNCB2467-100**

|                          |   |
|--------------------------|---|
| Artikelname              | Perforin-1 (Pore Forming Protein) (Apoptosis Marker)(PRF1/2467), Biotin conjugate, 0.1mg/mL, Clone: [PRF1/2467], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNCB2467-100  |
| Hersteller Artikelnummer | BNCB2467-100  |
| Alternativnummer         | BOT-BNCB2467-100-100UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant human Perforin-1 protein fragment (around aa 413-552) (exact sequence is proprietary)   |
| Konjugation              | Biotin  |
| Produktbeschreibung      | Perforin is a pore-forming protein that leads to osmotic lysis of the target cells and subsequently enables granzymes to enter the target cells and activate apoptosis. Perforin has structural and functional similarities to complement component 9 (C9)... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [PRF1/2467]   |

|                        |  |
|------------------------|--|
| Molekulargewicht       | 75 kDa   |
| UniProt                | <a href="#">P14222</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |