

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Bcl-6 (Follicular Lymphoma Marker) (rBCL6/1718), Biotin conjugate, 0.1mg/mL, Clone: [rBCL6/1718], Mouse, Monoclonal BOT-BNCB2498-100

Artikelname	Bcl-6 (Follicular Lymphoma Marker) (rBCL6/1718), Biotin conjugate, 0.1mg/mL, Clone: [rBCL6/1718], Mouse, Monoclonal
Artikelnummer	BOT-BNCB2498-100
Hersteller Artikelnummer	BNCB2498-100
Alternativnummer	BOT-BNCB2498-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human bcl-6 protein fragment (aa256-389) (exact sequence is proprietary)
Konjugation	Biotin
Produktbeschreibung	Bcl-6 is a transcriptional repressor mainly required for germinal center (GC) formation and antibody affinity maturation which has different mechanisms of action specific to the lineage and biological functions. It forms complexes with different core...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[rBCL6/1718]
Molekulargewicht	95 kDa
UniProt	P41182
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user