

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Galectin-1 / Human Placental Lactogen (hPL) (GAL1/2499R), Biotin conjugate, 0.1mg/mL, Clone: [GAL1/2499R], Rabbit, Monoclonal BOT-BNCB2499-500

| | |
|--------------------------|---|
| Artikelname | Galectin-1 / Human Placental Lactogen (hPL) (GAL1/2499R), Biotin conjugate, 0.1mg/mL, Clone: [GAL1/2499R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNCB2499-500 |
| Hersteller Artikelnummer | BNCB2499-500 |
| Alternativnummer | BOT-BNCB2499-500-500UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human Galectin-1 protein fragment (aa12-108) (exact sequence is proprietary) |
| Konjugation | Biotin |
| Produktbeschreibung | Galectin-1 is a member of the beta-galactoside-binding family and is a dimeric protein of 14 kD participating in a variety of normal and pathological processes, including cancer progression. Galectin-1 can affect the proliferation of normal and malig... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [GAL1/2499R] |
| Molekulargewicht | 14 kDa |
| UniProt | P09382 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody ELISA: 2-4 ug/m for coating order Ab without BSA Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |