

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD68 (Macrophage Marker) (C68/2501), Biotin conjugate, 0.1mg/mL, Clone: [C68/2501], Mouse, Monoclonal BOT-BNCB2501-500

Artikelname	CD68 (Macrophage Marker) (C68/2501), Biotin conjugate, 0.1mg/mL, Clone: [C68/2501], Mouse, Monoclonal
Artikelnummer	BOT-BNCB2501-500
Hersteller Artikelnummer	BNCB2501-500
Alternativnummer	BOT-BNCB2501-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant fragment of human CD68 protein (around aa 150-221) (exact sequence is proprietary)
Konjugation	Biotin
Produktbeschreibung	This antibody recognizes a glycoprotein of 110 kDa, which is identified as CD68. It is important for identifying macrophages in tissue sections. It stains macrophages in a wide variety of human tissues, including Kupffer cells and macrophages in the ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[C68/2501]
Molekulargewicht	~110 kDa
UniProt	P34810
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Flow cytometry: 0.5-1 ug/million cells Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 minutes followed by cooling at RT for 20 minutes Western: 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user