

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

DOG-1 / TMEM16A (Gastrointestinal Stromal Tumor Marker) (DG1/2564R), Biotin conjugate, 0.1mg/mL, Clone: [DG1/2564R], Rabbit, Monoclonal BOT-BNCB2564-500

| | |
|--------------------------|---|
| Artikelname | DOG-1 / TMEM16A (Gastrointestinal Stromal Tumor Marker) (DG1/2564R), Biotin conjugate, 0.1mg/mL, Clone: [DG1/2564R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNCB2564-500 |
| Hersteller Artikelnummer | BNCB2564-500 |
| Alternativnummer | BOT-BNCB2564-500-500UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human DOG-1 protein fragment (aa 2-101) (exact sequence is proprietary) |
| Konjugation | Biotin |
| Produktbeschreibung | Expression of DOG-1 protein is elevated in the gastrointestinal stromal tumors (GIST s), c-kit signaling-driven mesenchymal tumors of the GI tract. DOG-1 is rarely expressed in other soft tissue tumors, which, due to appearance, may be difficult to d... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [DG1/2564R] |
| Molekulargewicht | ~114 kDa |
| UniProt | Q5XXA6 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |