

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD3e (T-Cell Marker)(rC3e/1308), Biotin conjugate, 0.1mg/mL, Clone: [rC3e/1308], Mouse, Monoclonal BOT-BNCB3631-100**

Artikelname	CD3e (T-Cell Marker)(rC3e/1308), Biotin conjugate, 0.1mg/mL, Clone: [rC3e/1308], Mouse, Monoclonal
Artikelnummer	BOT-BNCB3631-100
Hersteller Artikelnummer	BNCB3631-100
Alternativnummer	BOT-BNCB3631-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant full-length human CD3E protein
Konjugation	Biotin
Produktbeschreibung	Recognizes the epsilon-chain of CD3, which consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28 kDa. The CD3 complex is closely associated at the lymphocyte cell surface with th...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[rC3e/1308]

Molekulargewicht	20 kDa
UniProt	<a href="#">P07766</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Not recommended for flow cytometry Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Western blotting 1-2 ug/mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user