

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ep-CAM / CD326(VU-1D9), 0.2mg/mL, Clone: [VU-1D9], Mouse, Monoclonal BOT-BNUB0017-500

Artikelname	Ep-CAM / CD326(VU-1D9), 0.2mg/mL, Clone: [VU-1D9], Mouse, Monoclonal
Artikelnummer	BOT-BNUB0017-500
Hersteller Artikelnummer	BNUB0017-500
Alternativnummer	BOT-BNUB0017-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	Exosome Staining, FC, IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Small cell lung carcinoma cells
Produktbeschreibung	This antibody reacts with the first EGF repeat in the extracellular domain of Ep-CAM. It is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[VU-1D9]
Molekulargewicht	40-43 kDa

UniProt	P16422
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with rat or ferret, others not known Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user