

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Plasma Cell Marker(LIV3G11, same as 7B18), 0.2mg/mL, IgG2a, Clone: [LIV3G11 same as 7B18], Mouse, Monoclonal BOT-BNUB0047-500**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Plasma Cell Marker(LIV3G11, same as 7B18), 0.2mg/mL, IgG2a, Clone: [LIV3G11 same as 7B18], Mouse, Monoclonal                                                                                                                                                  |
| Artikelnummer            | BOT-BNUB0047-500                                                                                                                                                                                                                                              |
| Hersteller Artikelnummer | BNUB0047-500                                                                                                                                                                                                                                                  |
| Alternativnummer         | BOT-BNUB0047-500-500UL                                                                                                                                                                                                                                        |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IF, IHC                                                                                                                                                                                                                                                       |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Pancreatic cancer-related mucin                                                                                                                                                                                                                               |
| Produktbeschreibung      | This antibody recognizes an intra-cytoplasmic antigen, which shows a very high degree of specificity for plasma cells. This antigen is present in normal as well as neoplastic plasma cells. Plasma cells, which are large lymphocytes derived from an ant... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.2 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [LIV3G11 same as 7B18]                                                                                                                                                                                                                                        |
| Molekulargewicht         | Not Known                                                                                                                                                                                                                                                     |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Isotyp                 | IgG2a                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| UniProt                | Not Known                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Does not react with rat, others not known Immunohistology formalin-fixed 0.1-0.2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user</p> |