

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

IgA Secretory Component(SC05), 0.2mg/mL, Clone: [SC05], Mouse, Monoclonal BOT-BNUB0050-100

Artikelname	IgA Secretory Component(SC05), 0.2mg/mL, Clone: [SC05], Mouse, Monoclonal
Artikelnummer	BOT-BNUB0050-100
Hersteller Artikelnummer	BNUB0050-100
Alternativnummer	BOT-BNUB0050-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	Functional Studies, IHC, WB
Spezies Reaktivität	Human, Rat
Immunogen	Secretory Component protein isolated from human colostrum
Produktbeschreibung	This MAb reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[SC05]
Molekulargewicht	~80 kDa

UniProt	Q16610
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user