

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD45 / LCA(136-4B5), 0.2mg/mL, Clone: [136-4B5], Mouse, Monoclonal BOT-BNUB0173-500

| | |
|--------------------------|---|
| Artikelname | CD45 / LCA(136-4B5), 0.2mg/mL, Clone: [136-4B5], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB0173-500 |
| Hersteller Artikelnummer | BNUB0173-500 |
| Alternativnummer | BOT-BNUB0173-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Stimulated human leukocytes |
| Produktbeschreibung | CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms ar... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [136-4B5] |
| Molekulargewicht | 180-220 kDa |

| | |
|------------------------|--|
| UniProt | P08575 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Flow cytometry: 1-2 ug/million cells Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |