

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD50 (ICAM-3)(101-1D2), 0.2mg/mL, Clone: [101-1D2], Mouse, Monoclonal BOT-BNUB0177-500

Artikelname	CD50 (ICAM-3)(101-1D2), 0.2mg/mL, Clone: [101-1D2], Mouse, Monoclonal
Artikelnummer	BOT-BNUB0177-500
Hersteller Artikelnummer	BNUB0177-500
Alternativnummer	BOT-BNUB0177-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	ELISA, FC, Functional Studies
Spezies Reaktivität	Human
Immunogen	Stimulated human leukocytes
Produktbeschreibung	Recognizes an N-glycosylated glycoprotein of 120 kDa with intra-chain disulfide bonds, identified as CD50 or ICAM-3 (WS: IV & V). Its epitope localizes in the D2 extracellular domain and is resistant to neuraminidase and proteases. CD50 is the major ...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[101-1D2]
Molekulargewicht	110-160 kDa

UniProt	P32942
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin/paraffin tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user