

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MHC II (HLA-DP) (beta chain)(BRA-FB6), 0.2mg/mL, Clone: [BRA-FB6], Mouse, Monoclonal BOT-BNUB0199-100

Artikelname	MHC II (HLA-DP) (beta chain)(BRA-FB6), 0.2mg/mL, Clone: [BRA-FB6], Mouse, Monoclonal
Artikelnummer	BOT-BNUB0199-100
Hersteller Artikelnummer	BNUB0199-100
Alternativnummer	BOT-BNUB0199-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	ELISA, FC, Functional Studies, IP
Spezies Reaktivität	Human
Immunogen	Non-T, non-B human acute lymphoblastic leukemia REH6 cell line
Produktbeschreibung	Recognizes a non-polymorphic determinant of DP-MHC class II. MHC class II antigens are transmembrane glycoproteins of non-covalently linked alpha (33-35 kDa) and beta (27-30 kDa) chains. It reportedly reacts with B- & non-T, non-B cell lines but not w...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[BRA-FB6]
Molekulargewicht	33-35 kDa

UniProt	P04440
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user