

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD15 / FUT4 / Lewis x(MMA (Leu-M1)), 0.2mg/mL, Clone: [Leu-M1], Mouse, Monoclonal BOT-BNUB0277-100

| | |
|--------------------------|---|
| Artikelname | CD15 / FUT4 / Lewis x(MMA (Leu-M1)), 0.2mg/mL, Clone: [Leu-M1], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB0277-100 |
| Hersteller Artikelnummer | BNUB0277-100 |
| Alternativnummer | BOT-BNUB0277-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | U937 histiocytic cell line |
| Produktbeschreibung | CD15 plays a role in mediating phagocytosis, bactericidal activity, and chemotaxis. It is present on >95% of granulocytes including neutrophils and eosinophils and to a lesser degree on monocytes. In addition, CD15 is expressed in Reed-Sternberg cell... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [Leu-M1] |
| Molekulargewicht | 220 kDa |

| | |
|------------------------|---|
| UniProt | P22083 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Immunofluorescence 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells Optimal dilution for a specific application should be determined by user |