

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Lewis A(7LE), 0.2mg/mL, Clone: [7LE], Mouse, Monoclonal BOT-BNUB0311-100

| | |
|--------------------------|---|
| Artikelname | Lewis A(7LE), 0.2mg/mL, Clone: [7LE], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB0311-100 |
| Hersteller Artikelnummer | BNUB0311-100 |
| Alternativnummer | BOT-BNUB0311-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | ELISA, FC, IHC |
| Spezies Reaktivität | Human, Mouse |
| Immunogen | Mucins isolated from ovarian cyst fluid |
| Produktbeschreibung | Recognizes a carbohydrate determinant of Gal 1-3(Fuc 1-4) GlcNAc which is blood group antigen Lewis A. It is present primarily on epithelial cells such as colon and kidneys. In the tumors and dedifferentiated tissues, decrease of Lewis A antigen was ... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [7LE] |
| Molekulargewicht | Multiple |
| UniProt | Not Applicable |

| | |
|------------------------|--|
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |