

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

BCL-10(BL10/411), 0.2mg/mL, Clone: [BL10/411], Mouse, Monoclonal BOT-BNUB0411-500

Artikelname	BCL-10(BL10/411), 0.2mg/mL, Clone: [BL10/411], Mouse, Monoclonal
Artikelnummer	BOT-BNUB0411-500
Hersteller Artikelnummer	BNUB0411-500
Alternativnummer	BOT-BNUB0411-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IHC, WB
Spezies Reaktivität	Human
Immunogen	Human BCL10 recombinant protein (epitope aa122-168)
Produktbeschreibung	B-cell lymphoma/leukemia 10 (BCL10) is a signaling protein that promotes apoptosis, pro-caspase-9 maturation, and activation of NF-kappa-B via NIK and IKK. BCL10 contains an N-terminal caspase recruitment domain (CARD), which is found in a number of ...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[BL10/411]
Molekulargewicht	33 kDa
UniProt	O95999

Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user