

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

EGFR(GFR450), 0.2mg/mL, Clone: [GFR450], Mouse, Monoclonal BOT-BNUB0450-500

| | |
|--------------------------|---|
| Artikelname | EGFR(GFR450), 0.2mg/mL, Clone: [GFR450], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB0450-500 |
| Hersteller Artikelnummer | BNUB0450-500 |
| Alternativnummer | BOT-BNUB0450-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant extracellular domain of human EGFR protein |
| Produktbeschreibung | This MAb recognizes a protein of 170 kDa, identified as EGFR. EGFR is type I receptor tyrosine kinase with sequence homology to erbB-1, -2, -3 -4 or HER-1, -2, -3 -4. It binds to Epidermal Growth Factor (EGF), Transforming Growth Factor-a (TGF-a), He... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [GFR450] |
| Molekulargewicht | ~170 kDa (wild type) and ~145 kDa (vIII variant) |
| UniProt | P00533 |

| | |
|------------------------|---|
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Immunohistology formalin-fixed 2-4 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Does not react with mouse, others not known Optimal dilution for a specific application should be determined by user |