

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

c-Myc(9E10.3), 0.2mg/mL, Clone: [9E10.3], Mouse, Monoclonal BOT-BNUB0596-100

| | |
|--------------------------|---|
| Artikelname | c-Myc(9E10.3), 0.2mg/mL, Clone: [9E10.3], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB0596-100 |
| Hersteller Artikelnummer | BNUB0596-100 |
| Alternativnummer | BOT-BNUB0596-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | A synthetic peptide, corresponding to aa 408-439 (AEEQKLISEEDLLRKRREQLKHKLEQLRNCA) from C-terminus of human c-myc, coupled to KLH. |
| Produktbeschreibung | The epitope of this antibody spans the Myc tag epitope, aa 410-419 (EQKLISEEDL), which is a specific portion of an alpha helical region of human c-myc protein. Because the epitope is located at the dimerization site of the protein, the antibody is no... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [9E10.3] |
| Molekulargewicht | 62-64 kDa |

| | |
|------------------------|--|
| UniProt | P01106 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Does not react with mouse, rat, or chicken, others not tested Optimal dilution for a specific application should be determined by user |