

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Mucin 1 / EMA / Episialin / CD227(VU-2G7), 0.2mg/mL, Clone: [VU-2G7], Mouse, Monoclonal BOT-BNUB0630-100**

Artikelname	Mucin 1 / EMA / Episialin / CD227(VU-2G7), 0.2mg/mL, Clone: [VU-2G7], Mouse, Monoclonal
Artikelnummer	BOT-BNUB0630-100
Hersteller Artikelnummer	BNUB0630-100
Alternativnummer	BOT-BNUB0630-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	ELISA, FC, IHC
Spezies Reaktivität	Human
Immunogen	Synthetic glycosylated MUC1 60mer tandem repeat NH <sub>2</sub> -(HGVTSA <sub>P</sub> DT(GalINAc)RPAPGSTAPPAHG) <sub>3</sub> -COOH, conjugated to bovine serum albumin
Produktbeschreibung	MAb VU-2G7 reacts with MUC1, a large transmembrane glycoprotein expressed on the ductal surface of normal glandular epithelia. The dominant epitope of VU-2G7 has not been established with epitope fingerprinting. The extracellular domain of MUC1 large...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[VU-2G7]

Molekulargewicht	265-400 kDa
UniProt	<a href="#">P15941</a>
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Immunohistology formalin-paraffin 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user