

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **p27 / KIP1(SX53G8), 0.2mg/mL, Clone: [SX53G8], Mouse, Monoclonal BOT-BNUB0669-500**

|                          |   |
|--------------------------|---|
| Artikelname              | p27 / KIP1(SX53G8), 0.2mg/mL, Clone: [SX53G8], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNUB0669-500  |
| Hersteller Artikelnummer | BNUB0669-500  |
| Alternativnummer         | BOT-BNUB0669-500-500UL  |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC, IP, WB   |
| Spezies Reaktivität      | Human, Monkey, Mouse, Rat   |
| Immunogen                | Purified GST-p27 fusion protein of human origin   |
| Produktbeschreibung      | This MAb recognizes a 27 kDa protein, identified as the p27Kip1, a cell cycle regulatory mitotic inhibitor. It is highly specific and shows no cross-reaction with other related mitotic inhibitors. In Western blotting of cell lysates from 7 human brea... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.2 mg/mL   |
| Klon-Bezeichnung         | [SX53G8]  |
| Molekulargewicht         | 25-26 kDa   |
| UniProt                  | <a href="#">P46527</a>  |

|                        |  |
|------------------------|--|
| Puffer                 | PBS, 0.05% BSA, 0.05% azide  |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |