

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 8(C-43), 0.2mg/mL, Clone: [C-43], Mouse, Monoclonal BOT-BNUB0688-100

Artikelname	Cytokeratin 8(C-43), 0.2mg/mL, Clone: [C-43], Mouse, Monoclonal
Artikelnummer	BOT-BNUB0688-100
Hersteller Artikelnummer	BNUB0688-100
Alternativnummer	BOT-BNUB0688-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC
Spezies Reaktivität	Bovine, Human, Porcine, Rabbit, Sheep
Immunogen	Cytoskeletal preparation from HeLa cells
Produktbeschreibung	Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenoc...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[C-43]
Molekulargewicht	52.5 kDa
UniProt	P05787

Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with mouse, rat, chicken or Xenopus Immunohistology formalin-fixed 0.5-1.0 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user