

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD57 / B3GAT1(NK-1 + HNK-1), 0.2mg/mL, IgM, Clone: [NK-1 HNK-1], Mouse, Monoclonal BOT-BNUB0695-500

Artikelname	CD57 / B3GAT1(NK-1 + HNK-1), 0.2mg/mL, IgM, Clone: [NK-1 HNK-1], Mouse, Monoclonal
Artikelnummer	BOT-BNUB0695-500
Hersteller Artikelnummer	BNUB0695-500
Alternativnummer	BOT-BNUB0695-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Human peripheral blood mononuclear cells (HNK-1 & NK-1)
Produktbeschreibung	Anti-CD57 marks a subset of lymphocytes known as natural killer (NK) cells. Follicular center cell lymphomas often contain many NK cells within the neoplastic follicles. Anti-CD57 also stains neuroendocrine cells and their derived tumors, including c...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[NK-1 HNK-1]
Molekulargewicht	~110 kDa

Isotyp	IgM
UniProt	Q9P2W7
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Does not react with rat, others not known Immunohistology formalin-fixed 2-4 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user