

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Melanoma Marker(HMB45 + M2-7C10 + M2-9E3 + T311), 0.2mg/mL, Clone: [HMB45 M2-7C10 M2-9E3 T311], Mouse, Monoclonal BOT-BNUB0701-500

| | |
|--------------------------|---|
| Artikelname | Melanoma Marker(HMB45 + M2-7C10 + M2-9E3 + T311), 0.2mg/mL, Clone: [HMB45 M2-7C10 M2-9E3 T311], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB0701-500 |
| Hersteller Artikelnummer | BNUB0701-500 |
| Alternativnummer | BOT-BNUB0701-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant hMART-1 protein (M2-7C10, M2-9E3), Recombinant tyrosinase protein (T311), Extract of pigmented melanoma metastases from lymph nodes (HMB45) |
| Produktbeschreibung | This antibody cocktail recognizes three melanoma-specific proteins, which include MART-1, Tyrosinase and gp100. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Tyrosinase is one of the... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [HMB45 M2-7C10 M2-9E3 T311] |

| | |
|------------------------|---|
| Molekulargewicht | 20-22 kDa (doublet) (MART), 70-80 kDa (Tyrosinase), 90-100 kDa (gp100) |
| UniProt | Q16655 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (Formalin-fixed) 0.5-1.0 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |