

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p53(BP53-12 + DO-7), 0.2mg/mL, Clone: [BP53-12 DO-7], Mouse, Monoclonal BOT-BNUB0723-100

Artikelname	p53(BP53-12 + DO-7), 0.2mg/mL, Clone: [BP53-12 DO-7], Mouse, Monoclonal
Artikelnummer	BOT-BNUB0723-100
Hersteller Artikelnummer	BNUB0723-100
Alternativnummer	BOT-BNUB0723-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Bovine, Human, Monkey
Immunogen	Recombinant human wild-type p53 protein (BP53-12) & Recombinant human wild type p53 protein expressed in E. coli (DO-7)
Produktbeschreibung	Recognizes a 53 kDa protein, which is identified as p53 suppressor gene product. It reacts with the mutant as well as the wild form of p53 under denaturing and non-denaturing conditions. Its epitope maps within the N-terminus (aa 20-25) of p53 oncopr...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[BP53-12 DO-7]

Molekulargewicht	53 kDa
UniProt	P04637
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user