

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p53(BP53-12 + DO-7), 0.2mg/mL, Clone: [BP53-12 DO-7], Mouse, Monoclonal BOT-BNUB0723-500

| | |
|--------------------------|---|
| Artikelname | p53(BP53-12 + DO-7), 0.2mg/mL, Clone: [BP53-12 DO-7], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB0723-500 |
| Hersteller Artikelnummer | BNUB0723-500 |
| Alternativnummer | BOT-BNUB0723-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Bovine, Human, Monkey |
| Immunogen | Recombinant human wild-type p53 protein (BP53-12) & Recombinant human wild type p53 protein expressed in E. coli (DO-7) |
| Produktbeschreibung | Recognizes a 53 kDa protein, which is identified as p53 suppressor gene product. It reacts with the mutant as well as the wild form of p53 under denaturing and non-denaturing conditions. Its epitope maps within the N-terminus (aa 20-25) of p53 oncopr... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [BP53-12 DO-7] |

| | |
|------------------------|--|
| Molekulargewicht | 53 kDa |
| UniProt | P04637 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user |