

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Thyroglobulin(2H11 + 6E1), 0.2mg/mL, Clone: [2H11 6E1], Mouse, Monoclonal BOT-BNUB0724-100**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Thyroglobulin(2H11 + 6E1), 0.2mg/mL, Clone: [2H11 6E1], Mouse, Monoclonal                                                                                                                                                                                     |
| Artikelnummer            | BOT-BNUB0724-100                                                                                                                                                                                                                                              |
| Hersteller Artikelnummer | BNUB0724-100                                                                                                                                                                                                                                                  |
| Alternativnummer         | BOT-BNUB0724-100-100UL                                                                                                                                                                                                                                        |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC                                                                                                                                                                                                                                                           |
| Spezies Reaktivität      | Human, Mouse, Rat                                                                                                                                                                                                                                             |
| Immunogen                | Human thyroid follicular cells                                                                                                                                                                                                                                |
| Produktbeschreibung      | Thyroglobulin is a 660 kDa dimeric pre-protein with mutiple glycosylation sites. It is produced by and processed within the thyroid gland to produce the hormone thyroxine and triiodothyronine. Prior to forming dimers, thyroglobulin monomers undergo c... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.2 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [2H11 6E1]                                                                                                                                                                                                                                                    |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Molekulargewicht       | 660 kDa (Dimeric Form)                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| UniProt                | <a href="#">P01266</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology formalin-fixed 0.1-0.2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1.0 ug/million cells in 0.1 mL Optimal dilution for a specific application should be determined by user</p> |