

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human Lambda Light Chain(LcN-2 + ICO-106), 0.2mg/mL, Clone: [LcN-2 ICO-106], Mouse, Monoclonal BOT-BNUB0735-100

| | |
|--------------------------|---|
| Artikelname | Human Lambda Light Chain(LcN-2 + ICO-106), 0.2mg/mL, Clone: [LcN-2 ICO-106], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB0735-100 |
| Hersteller Artikelnummer | BNUB0735-100 |
| Alternativnummer | BOT-BNUB0735-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | WB |
| Spezies Reaktivität | Human |
| Immunogen | Purified human IgG (LcN-2 and ICO-106) |
| Produktbeschreibung | This MAb is specific to lambda light chain of immunoglobulin and shows no cross-reaction with lambda light chain or any of the five heavy chains. In mammals, the two light chains in an antibody are always identical, with only one type of light chain,... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [LcN-2 ICO-106] |
| Molekulargewicht | ~22.5 kDa |

| | |
|------------------------|--|
| UniProt | P01701 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user |