

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

bcl-x(BX006 + 2H12), 0.2mg/mL, Clone: [BX006 2H12], Mouse, Monoclonal BOT-BNUB0752-500

Artikelname	bcl-x(BX006 + 2H12), 0.2mg/mL, Clone: [BX006 2H12], Mouse, Monoclonal
Artikelnummer	BOT-BNUB0752-500
Hersteller Artikelnummer	BNUB0752-500
Alternativnummer	BOT-BNUB0752-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human, Mouse, Porcine, Rat
Immunogen	A synthetic peptide, aa 3-14 (Cys-QSNRELVVDFLS) of human Bcl-X protein (BX006 & 2H12)
Produktbeschreibung	Bcl-x belongs to the Bcl-2 family of proteins and plays a crucial role in apoptosis. Bcl-X has two isoforms: Bcl-XL (long), a 26 kDa protein which suppresses cell death, and Bcl-XS (short), a 178 amino acid protein lacking a 63 amino acid domain whic...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[BX006 2H12]
Molekulargewicht	27 kDa

UniProt	Q07817
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user