

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 19(A53-B/A2.26 + BA17), 0.2mg/mL, Clone: [A53-B/A2.26 + BA17], Mouse, Monoclonal BOT-BNUB0753-100

Artikelname	Cytokeratin 19(A53-B/A2.26 + BA17), 0.2mg/mL, Clone: [A53-B/A2.26 + BA17], Mouse, Monoclonal
Artikelnummer	BOT-BNUB0753-100
Hersteller Artikelnummer	BNUB0753-100
Alternativnummer	BOT-BNUB0753-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Human breast cancer MCF-7 cells (A53-B/A2), Human mammary epithelial organoids (BA17)
Produktbeschreibung	This MAb reacts with the rod domain of human cytokeratin-19 (CK19), a polypeptide of 40 kDa. CK19 is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[A53-B/A2.26 + BA17]

Molekulargewicht	40 kDa
UniProt	P08727
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user