

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p27 / KIP1(DCS-72.F6), 0.2mg/mL, Clone: [DCS-72.F6], Mouse, Monoclonal BOT-BNUB0771-500

| | |
|--------------------------|---|
| Artikelname | p27 / KIP1(DCS-72.F6), 0.2mg/mL, Clone: [DCS-72.F6], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB0771-500 |
| Hersteller Artikelnummer | BNUB0771-500 |
| Alternativnummer | BOT-BNUB0771-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IHC |
| Spezies Reaktivität | Human, Monkey, Mouse, Rat |
| Immunogen | Mouse recombinant p27 protein |
| Produktbeschreibung | Recognizes a 27 kDa protein, identified as the p27Kip1, a cell cycle regulatory mitotic inhibitor. Its epitope spans between aa 83-204 of p27. It is highly specific and shows no cross-reaction with other related mitotic inhibitors. p27Kip1 functions ... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [DCS-72.F6] |
| Molekulargewicht | 25-26 kDa |

| | |
|------------------------|--|
| UniProt | P46527 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |