

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 19(KRT19/799), 0.2mg/mL, Clone: [KRT19/799], Mouse, Monoclonal BOT-BNUB0799-500

| | |
|--------------------------|---|
| Artikelname | Cytokeratin 19(KRT19/799), 0.2mg/mL, Clone: [KRT19/799], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB0799-500 |
| Hersteller Artikelnummer | BNUB0799-500 |
| Alternativnummer | BOT-BNUB0799-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human KRT19 protein |
| Produktbeschreibung | Recognizes a protein of 40 kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epit... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [KRT19/799] |
| Molekulargewicht | 40 kDa |

| | |
|------------------------|---|
| UniProt | P08727 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Methanol fixation recommended for immunofluorescence on cells Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user |