

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cyclin D1(CCND1/809), 0.2mg/mL, Clone: [CCND1/809], Mouse, Monoclonal BOT-BNUB0809-500

| | |
|--------------------------|---|
| Artikelname | Cyclin D1(CCND1/809), 0.2mg/mL, Clone: [CCND1/809], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB0809-500 |
| Hersteller Artikelnummer | BNUB0809-500 |
| Alternativnummer | BOT-BNUB0809-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human CCND1 protein |
| Produktbeschreibung | Recognizes a protein of 36 kDa, identified as cyclin D1. Cyclin D1, one of the key cell cycle regulators, is a putative proto-oncogene overexpressed in a wide variety of human neoplasms. This antibody neutralizes the activity of cyclin D1 in vivo. Ab... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [CCND1/809] |
| Molekulargewicht | 36 kDa |

| | |
|------------------------|---|
| UniProt | P24385 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |