

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Involucrin(IVRN/827), 0.2mg/mL, Clone: [IVRN/827], Mouse, Monoclonal BOT-BNUB0827-100

| | |
|--------------------------|---|
| Artikelname | Involucrin(IVRN/827), 0.2mg/mL, Clone: [IVRN/827], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB0827-100 |
| Hersteller Artikelnummer | BNUB0827-100 |
| Alternativnummer | BOT-BNUB0827-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Purified involucrin from human keratinocytes |
| Produktbeschreibung | This antibody recognizes a protein of 66 kDa-170 kDa, identified as involucrin. In Western blotting of cultured human keratinocytes, this MAb reacts with a 120 kDa protein. Involucrin is expressed in a range of stratified squamous epithelia, includin... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [IVRN/827] |
| Molekulargewicht | 66-170 kDa |

| | |
|------------------------|---|
| UniProt | P07476 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.1-0.2 ug/mL Staining of formalin-fixed tissues is enhanced by digestion with trypsin or Protease XXV at 1mg/mL PBS for 5 min at 37C Flow Cytometry 0.5-1 ug/million cells/0.1 mL Note than enzyme digestion is better than citrate buffer based epitope unmasking Optimal dilution for a specific application should be determined by user |