

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mucin 1 / EMA / Episialin / CD227(MUC1/845), 0.2mg/mL, Clone: [MUC1/845], Mouse, Monoclonal BOT-BNUB0845-100

Artikelname	Mucin 1 / EMA / Episialin / CD227(MUC1/845), 0.2mg/mL, Clone: [MUC1/845], Mouse, Monoclonal
Artikelnummer	BOT-BNUB0845-100
Hersteller Artikelnummer	BNUB0845-100
Alternativnummer	BOT-BNUB0845-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Human milk-fat globule membranes (HMFGM)
Produktbeschreibung	In Western blotting, it recognizes proteins in MW range of 265-400 kDa, identified as different glycoforms of EMA. The alpha subunit has cell adhesive properties. It can act both as an adhesion and an anti-adhesion protein. EMA may provide a protectiv...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[MUC1/845]
Molekulargewicht	265-400 kDa

UniProt	P15941
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.1-0.2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user