

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TTF-1 / NKX2.1(8G7G3/1 + NX2.1/690), 0.2mg/mL, Clone: [8G7G3/1 NX2.1/690], Mouse, Monoclonal BOT-BNUB0907-100

Artikelname	TTF-1 / NKX2.1(8G7G3/1 + NX2.1/690), 0.2mg/mL, Clone: [8G7G3/1 NX2.1/690], Mouse, Monoclonal
Artikelnummer	BOT-BNUB0907-100
Hersteller Artikelnummer	BNUB0907-100
Alternativnummer	BOT-BNUB0907-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Rat full length TTF-1 recombinant protein (8G7G3/1), Recombinant TTF-1 protein (NX2.1/690)
Produktbeschreibung	Recognizes a protein of 40 kDa, identified as Thyroid transcription factor-1 (TTF-1). TTF-1 is a member of the NKx2 family of homeodomain transcription factors. It is expressed in epithelial cells of the thyroid gland and the lung. Nuclei from liver,...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[8G7G3/1 NX2.1/690]

Molekulargewicht	40 kDa
UniProt	P43699
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user