

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Mucin 1 / EMA / Episialin / CD227(139H2), 0.2mg/mL, Clone: [139H2], Mouse, Monoclonal BOT-BNUB0925-100**

Artikelname	Mucin 1 / EMA / Episialin / CD227(139H2), 0.2mg/mL, Clone: [139H2], Mouse, Monoclonal
Artikelnummer	BOT-BNUB0925-100
Hersteller Artikelnummer	BNUB0925-100
Alternativnummer	BOT-BNUB0925-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human, Mouse
Immunogen	Human milk-fat globule membranes (HMFGM)
Produktbeschreibung	This MAb reacts with MUC1. The dominant epitope of this MAb has not yet been determined. MUC1 is a large cell surface mucin glycoprotein expressed by most glandular and ductal epithelial cells and some hematopoietic cell lineages. It is expressed on ...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[139H2]
Molekulargewicht	265-400 kDa

UniProt	<a href="#">P15941</a>
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes ELISA For coating use Ab at 2-5 ug/mL, order Ab without BSA Flow Cytometry 0.25-0.5 ug/million cells in 0.1 mL Optimal dilution for a specific application should be determined by user