

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Thomsen-Friedenreich Antigen / CD176 (Pan Carcinoma Marker)(A63-C/A9), 0.2mg/mL, Clone: [A63-C/A9], Mouse, Monoclonal BOT-BNUB0938-100**

Artikelname	Thomsen-Friedenreich Antigen / CD176 (Pan Carcinoma Marker)(A63-C/A9), 0.2mg/mL, Clone: [A63-C/A9], Mouse, Monoclonal
Artikelnummer	BOT-BNUB0938-100
Hersteller Artikelnummer	BNUB0938-100
Alternativnummer	BOT-BNUB0938-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Neuraminidase-treated human red blood cells
Produktbeschreibung	This antibody recognizes a disaccharide epitope, Gal 1-3GalNAc, of Thomsen-Friedenreich (TF) antigen. It is specific for both anomeric forms of the disaccharide (TF and TF , including related structures on the glycolipid) and shows no cross-reactivit...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[A63-C/A9]
Molekulargewicht	Multiple

UniProt	Not Applicable
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at 37°C Immunohistochemistry: frozen: 0.5-1.0 ug/mL for 30 minutes at RT Immunofluorescence: 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user