

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD147(BSG/963), 0.2mg/mL, IgG1, Clone: [BSG/963], Mouse, Monoclonal BOT-BNUB0963-500

| | |
|--------------------------|---|
| Artikelname | CD147(BSG/963), 0.2mg/mL, IgG1, Clone: [BSG/963], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB0963-500 |
| Hersteller Artikelnummer | BNUB0963-500 |
| Alternativnummer | BOT-BNUB0963-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human BSG protein |
| Produktbeschreibung | This MAb recognizes extracellular epitope of human CD147. It is expressed more intensely on thymocytes than on mature peripheral blood T cells. CD147 is important in spermatogenesis, embryo implantation, neural network formation, and tumor progressio... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [BSG/963] |
| Molekulargewicht | 35 kDa (non-reduced), 40 kDa (reduced) |

| | |
|------------------------|---|
| Isotyp | IgG1 |
| UniProt | P35613 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Immunofluorescence 1:100-1:200 Optimal dilution for a specific application should be determined by user |