

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Ep-CAM / CD326(Cocktail PAN-EpCAM), 0.2mg/mL, Clone: [Cocktail PAN-EpCAM], Mouse, Monoclonal BOT-BNUB0969-500**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | Ep-CAM / CD326(Cocktail PAN-EpCAM), 0.2mg/mL, Clone: [Cocktail PAN-EpCAM], Mouse, Monoclonal                                                                                                                                                                  |
| Artikelnummer            | BOT-BNUB0969-500                                                                                                                                                                                                                                              |
| Hersteller Artikelnummer | BNUB0969-500                                                                                                                                                                                                                                                  |
| Alternativnummer         | BOT-BNUB0969-500-500UL                                                                                                                                                                                                                                        |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Mouse                                                                                                                                                                                                                                                         |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | Exosome Staining, FC, IF, IHC, WB                                                                                                                                                                                                                             |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant human Ep-CAM protein (full-length and fragments)                                                                                                                                                                                                  |
| Produktbeschreibung      | This is a cocktail of four highly specific monoclonal antibodies (EGP40/826, EGP40/837, EGP40/1110, EGP40/1120) that recognize extracellular as well as intracellular domains of the epithelial cellular adhesion molecule (Ep-CAM). Epithelial specific a... |
| Klonalität               | Monoclonal                                                                                                                                                                                                                                                    |
| Konzentration            | 0.2 mg/mL                                                                                                                                                                                                                                                     |
| Klon-Bezeichnung         | [Cocktail PAN-EpCAM]                                                                                                                                                                                                                                          |
| Molekulargewicht         | 40-43 kDa                                                                                                                                                                                                                                                     |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| UniProt                | <a href="#">P16422</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Puffer                 | PBS, 0.05% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |