

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Thymidine Phosphorylase / PD-ECGF (Angiogenesis Marker) (P-GF.44C), 0.2mg/mL, IgG1, Clone: [P-GF.44C], Mouse, Monoclonal BOT-BNUB1244-100

Artikelname	Thymidine Phosphorylase / PD-ECGF (Angiogenesis Marker) (P-GF.44C), 0.2mg/mL, IgG1, Clone: [P-GF.44C], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1244-100
Hersteller Artikelnummer	BNUB1244-100
Alternativnummer	BOT-BNUB1244-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Recombinant full-length human Thymidine Phosphorylase (TP / PD-ECGF) protein
Produktbeschreibung	This antibody recognizes a protein (amino acid 482) of 55 kDa (in vivo 110 kDa homodimer), identified as platelet-derived endothelial growth factor (PD-ECGF), same as thymidine phosphorylase (TP) or gliostatin. In the presence of inorganic orthophosp...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[P-GF.44C]

Molekulargewicht	55 kDa
Isotyp	IgG1
UniProt	P19971
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin) 1-2 ug/mL Western blotting 0.5-1 ug/mL Immunoprecipitation 0.5-1 ug/500ug protein lysate Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user