

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Napsin A (Lung Adenocarcinoma Marker)(NAPSA/1238 + NAPSA/1239), 0.2mg/mL, Clone: [NAPSA/1238 NAPSA/1239], Mouse, Monoclonal BOT-BNUB1254-500**

Artikelname	Napsin A (Lung Adenocarcinoma Marker)(NAPSA/1238 + NAPSA/1239), 0.2mg/mL, Clone: [NAPSA/1238 NAPSA/1239], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1254-500
Hersteller Artikelnummer	BNUB1254-500
Alternativnummer	BOT-BNUB1254-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human Napsin-A protein fragment (aa189-299) (exact sequence is proprietary)
Produktbeschreibung	Napsin is a pepsin-like aspartic proteinase connected with maturation of surfactant protein B. There are two closely related napsins, napsin A and napsin B. Napsin A is expressed as a single chain protein. Immunohistochemical studies revealed high exp...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[NAPSA/1238 NAPSA/1239]

Molekulargewicht	37 kDa
UniProt	<a href="#">O96009</a>
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT Western blot: 1-2 ug/mL for 60 minutes at RT Flow cytometry: 0.5-1 ug/million cells Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user