

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Human Immunoglobulin Gamma (IgG) Heavy Chain(IG217 + IG266), 0.2mg/mL, Clone: [IG217 IG266], Mouse, Monoclonal BOT-BNUB1262-100**

Artikelname	Human Immunoglobulin Gamma (IgG) Heavy Chain(IG217 + IG266), 0.2mg/mL, Clone: [IG217 IG266], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1262-100
Hersteller Artikelnummer	BNUB1262-100
Alternativnummer	BOT-BNUB1262-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Purified human IgG
Produktbeschreibung	This MAb is specific to heavy chain of IgG and shows minimal cross-reaction with heavy chains of other immunoglobulins. It is reactive with all subclasses of Gamma heavy chain(IgG1, IgG2a, IgG2b, IgG3). The most common feature of plasmacytomas, and c...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[IG217 IG266]
Molekulargewicht	75 kDa

UniProt	<a href="#">P01857</a>
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Flow cytometry: 1 ug/million cells Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT Immunofluorescence: 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user