

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

c-Myc Oncoprotein(MYC275 + MYC909), 0.2mg/mL, Clone: [MYC275 MYC909], Mouse, Monoclonal BOT-BNUB1269-500

| | |
|--------------------------|---|
| Artikelname | c-Myc Oncoprotein(MYC275 + MYC909), 0.2mg/mL, Clone: [MYC275 MYC909], Mouse, Monoclonal |
| Artikelnummer | BOT-BNUB1269-500 |
| Hersteller Artikelnummer | BNUB1269-500 |
| Alternativnummer | BOT-BNUB1269-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human c-myc protein |
| Produktbeschreibung | This antibody recognizes a transcription factor of 64-67 kDa, identified as c-myc. This MAb shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or over-expressed in a var... |
| Klonalität | Monoclonal |
| Konzentration | 0.2 mg/mL |
| Klon-Bezeichnung | [MYC275 MYC909] |
| Molekulargewicht | 62-64 kDa |

| | |
|------------------------|--|
| UniProt | P01106 |
| Puffer | PBS, 0.05% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistochemistry (formalin-fixed): 1-2 ug/mL for 30 minutes at RT Flow cytometry: 0.5-1 ug/million cells Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |