

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Human Kappa Light Chain (B-Cell Marker) (KLC1278), 0.2mg/mL, Clone: [KLC1278], Mouse, Monoclonal BOT-BNUB1278-500**

Artikelname	Human Kappa Light Chain (B-Cell Marker) (KLC1278), 0.2mg/mL, Clone: [KLC1278], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1278-500
Hersteller Artikelnummer	BNUB1278-500
Alternativnummer	BOT-BNUB1278-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human full-length Ig kappa chain
Produktbeschreibung	This MAb is specific to kappa light chain of immunoglobulin and shows no cross-reaction with lambda light chain or any of the five heavy chains. It recognizes human Ig kappa light chains of both secreted and cell surface immunoglobulin. It detects al...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[KLC1278]
Molekulargewicht	~22.5 kDa

UniProt	<a href="#">P01601</a>
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 8.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user