

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **VEGI (Vascular Endothelial Growth Inhibitor) (VEGI/1283), 0.2mg/mL, Clone: [VEGI/1283], Mouse, Monoclonal BOT-BNUB1283-500**

Artikelname	VEGI (Vascular Endothelial Growth Inhibitor) (VEGI/1283), 0.2mg/mL, Clone: [VEGI/1283], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1283-500
Hersteller Artikelnummer	BNUB1283-500
Alternativnummer	BOT-BNUB1283-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human full-length VEGI fragment
Produktbeschreibung	VEGI is an anti-angiogenic cytokine that belongs to tumor necrosis factor superfamily, member 15 (TNFSF15). This protein is abundantly expressed in endothelial cells, but is not expressed in either B or T cells. The expression of this protein is indu...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[VEGI/1283]
Molekulargewicht	22 kDa

UniProt	<a href="#">O95150</a>
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Flow cytometry: 0.5-1 ug/million cells in 0.1mL Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Western blotting 1-2 ug/mL Optimal dilution for a specific application should be determined by user