

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD3e (T-Cell Marker) (C3e/1308), 0.2mg/mL, Clone: [C3e/1308], Mouse, Monoclonal BOT-BNUB1308-100**

Artikelname	CD3e (T-Cell Marker) (C3e/1308), 0.2mg/mL, Clone: [C3e/1308], Mouse, Monoclonal
Artikelnummer	BOT-BNUB1308-100
Hersteller Artikelnummer	BNUB1308-100
Alternativnummer	BOT-BNUB1308-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human CD3e extracellular domain fragment (aa 23-119)
Produktbeschreibung	The immunogen for this Mab is derived from the extracellular domain of CD3 epsilon chain. CD3 consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28 kDa. The CD3 complex is closel...
Klonalität	Monoclonal
Konzentration	0.2 mg/mL
Klon-Bezeichnung	[C3e/1308]

Molekulargewicht	20 kDa
UniProt	<a href="#">P07766</a>
Puffer	PBS, 0.05% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Not recommended for flow cytometry Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Western blotting 1-2 ug/mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user